

Product Information

Pirinase Allergy 0.05% Nasal Spray (fluticasone propionate 50 micrograms per spray). **Uses:** Prevention and treatment of allergic rhinitis. **Dosage and method of use:** Adults: Two sprays into each nostril once or twice a day if required. Maintenance: one spray per nostril once a day if possible. Use lowest possible maintenance dose. Max 4 sprays a day in each nostril. Children under 18 years: Not to be used. **Contraindications:** Hypersensitivity to ingredients. **Precautions:** Seek medical advice if inadequate relief after 7 days, or require more than 3 months continuous use. Seek medical advice before use with: concomitant corticosteroid use, fever, nasal/sinus infection, recent nasal injury/surgery, nasal ulceration, pregnancy, breast feeding. Caution due to interactions with cytochrome P450 3A4 inhibitors like ketoconazole and protease inhibitors, such as ritonavir and cobicistat which may increase systemic exposure to fluticasone. Caution commencing in those withdrawing from systemic steroids, particularly if impairment of adrenal function suspected. Systemic effects of nasal corticosteroids may occur, particularly at high doses prescribed for prolonged periods. May cause visual disturbance. Contains Benzalkonium Chloride which may cause bronchospasm. **Side effects:** Dryness and irritation of the nose and throat, unpleasant taste and smell, headache and epistaxis. Hypersensitivity reactions including skin rash and oedema of the face or tongue. Eye disorders such as glaucoma and blurred vision. Rarely anaphylaxis/ anaphylactic reactions and bronchospasm. See SPC for full details. **Legal category:** P. **PL number:** PL 44673/0099. **Product licence holder:** GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. **Package quantity and RSP (excl. VAT):** 60 sprays £8.69. **Date of last revision:** January 2018.